



For research use only

ISO9001

10x TBE Buffer

Product	Volume	Cat. No.
10x TBE Buffer	1 L	EBA-1047

Description.

Elpis Biotech's 10x TBE (Tris-Borate-EDTA Buffer) is used for polyacrylamide and agarose gel electrophoresis. TBE was originally used at a working strength of 1x for polyacrylamide gel and 0.5x for agarose gel electrophoresis. Premixed electrophoresis buffers eliminate variables which can distort data and give unreliable results.

10x Composition

890 mM Tris base
890 mM Boric acid
20 mM EDTA
pH at 25°C (1x): 8.2-8.4

Recommended Storage Condition

Store at room temperature (22-25°C)

Usage Information (applications)

Electrophoresis buffer for polyacrylamide and agarose gel electrophoresis.

Caution

Not for diagnostic use. The safety and efficacy of this product in diagnostic or other clinical uses has not been established.



For research use only

ISO9001

10x TBE Buffer

Product	Volume	Cat. No.
10x TBE Buffer	1 L	EBA-1047

Description.

Elpis Biotech's 10x TBE (Tris-Borate-EDTA Buffer) is used for polyacrylamide and agarose gel electrophoresis. TBE was originally used at a working strength of 1x for polyacrylamide gel and 0.5x for agarose gel electrophoresis. Premixed electrophoresis buffers eliminate variables which can distort data and give unreliable results.

10x Composition

890 mM Tris base
890 mM Boric acid
20 mM EDTA
pH at 25°C (1x): 8.2-8.4

Recommended Storage Condition

Store at room temperature (22-25°C)

Usage Information (applications)

Electrophoresis buffer for polyacrylamide and agarose gel electrophoresis.

Caution

Not for diagnostic use. The safety and efficacy of this product in diagnostic or other clinical uses has not been established.