



For research use only

ISO9001

# 1Kbp plus 100bp DNA Ladder Marker, Ready-to-use

Product	Conc.	Cat. No.	Remarks
1Kbp plus 100bp DNA Ladder Marker, Ready-to-use	1 ml (83 µg/ml)	EBM-1003	Ready-to-use

### Description

The 1Kbp plus 100bp DNA ladder marker is a mixture of double-stranded DNA fragments for broad range determination of the exact size of DNA fragments. The 1Kbp plus 100bp DNA ladder marker consists of total 18 DNA fragments ranging in size from 100 to 10,000 bp. Ruler bands (500, 1,000, 1,500, 5,000 bp) are more brighter than the other bands and helpful for easy size discrimination on the gel electrophoresis. The 1Kbp plus 100bp DNA ladder marker is supplied in a ready-to-use format. This ladder marker can be stained by ethidium bromide or any other known DNA staining methods.

### Storage Buffer

- Marker DNA : 83 µg in 1 ml of 10 mM Tris-HCl, pH8.0, 1 mM EDTA, 5% Glycerol, 0.005% Bromophenol Blue, and 0.005% Xylene Cyanol

### Recommended Storage Condition

- -20°C for 2 year
- 4°C for 6 months
- Room temperature (20-25°C) for 2 months


### Usage Information

- Concentration : 415 ng/5 µl (83 µg/ml)
- Recommended loading : 5-10 µl (100-200 lanes, ready-to-use)
- Range : 100 – 10,000 bp
- Number of bands : 18

### Cautions


- Always use the fresh tip to take out marker solution.  
(If you do not, trace amount of contaminated DNases from buffer tank may degrade marker DNA rapidly)
- Don't boil the product.
- Use appropriate % of gels for separation of 100 to 10,000 bp sizes  
(0.7 to 1.5% agarose gel is recommended)
- Confirm that the concentration of DNA staining dye is optimal before use.  
(Breakage or suboptimal concentration of ethidium bromide in gel is a main cause of low estimation of marker concentration or your DNA. 5 ng of DNA should be seen in normal condition)
- Loading volume and concentration should be optimized by gel size, well size, and running length.
- Low sized DNA bands can be gradually disappeared as running is progressing.  
(This is because some DNA is getting out from gel to buffer during horizontal electrophoresis, not because the DNA concentration is incorrect. This will be the same for your DNA)

1Kbp plus 100bp DNA Ladder Marker



5 µl/415 ng/lane ;  
1.5% agarose in 0.5x TAE, stained with ethidium bromide

### Migration Patterns in Different % of Agarose Gels



### Recommended Gel Percentages for Separation of Linear DNA

Agarose Gel, %	Range of Separation, bp	Polyacrylamide Gel, %	Range of Separation, bp
0.5	1,000 - 30,000	3.5	100 - 1,000
0.7	800 - 12,000	5	80 - 500
1	500 - 10,000	8	60 - 400
1.2	400 - 7,000	12	40 - 200
1.4	200 - 4,000	20	5 - 100
2	50 - 2,000		

### DNA Size Migration with Sample Loading Dyes

Agarose Concentration, %	Xylene cyanol FF	Bromophenol blue	Orange G
0.7 - 1.7	~4000 bp	~300 bp	~50 bp
2.5 - 3.0	~800 bp	~100 bp	~30 bp

### Composition of Gel Electrophoresis Buffers

Buffer	Working Concentration	Stock Concentration (per Liter)		
Tris-acetate (TAE)	1x	20 mM Tris-acetate	Tris base	96.9 g
		1 mM EDTA	Glacial acetic acid	22.84 ml
			0.5 M EDTA (pH8.0)	40 ml
Tris-borate (TBE)	0.5x	45 mM Tris-borate	Tris base	108 g
		1 mM EDTA	Boric acid	55 g
			0.5 M EDTA (pH8.0)	40 ml