




For research use only

ISO9001

Aat II

Product	Quantity	Cat. No.	Remarks
Aat II	250 unit	EBR-1001	5 unit/ μ l

Recognition site**Source***Acetobacter aceti.***Reaction Conditions**

Buffer 4. 20 mM Tris-acetate (pH7.9), 50 mM potassium acetate, 10 mM magnesium acetate, 1 mM DTT. Incubate at 37°C.

Concentration & Storage Condition

Store at -20°C. 5 unit/ μ l in 10 mM Tris-HCl (pH 7.4), 50 mM KCl, 1 mM DTT, 0.1 mM EDTA, 200 μ g/ml BSA, 50% (v/v) glycerol.

Heat Inactivation Condition

65°C for 20min.

Enzyme dilution

Dilute with Storage Buffer.

QC Tests

Activity, ligation and recutting, exo and endonuclease activity test, SDS-PAGE purity, performance tests.

Methylase Sensitivity

Blocked by CpG methylation (Cleavage of mammalian genomic DNA is blocked).

Buf4	37°C	65°C	CpG	BW
<small>BUFFER</small>	<small>REACTION</small>	<small>INACTIVATION</small>	<small>METHYL BLOCK</small>	<small>BLUE/WHITE</small>

Buffer	1	2	3	4
Activity	0	50	50	100




For research use only

ISO9001

Aat II

Product	Quantity	Cat. No.	Remarks
Aat II	250 unit	EBR-1001	5 unit/ μ l

Recognition site**Source***Acetobacter aceti.***Reaction Conditions**

Buffer 4. 20 mM Tris-acetate (pH7.9), 50 mM potassium acetate, 10 mM magnesium acetate, 1 mM DTT. Incubate at 37°C.

Concentration & Storage Condition

Store at -20°C. 5 unit/ μ l in 10 mM Tris-HCl (pH 7.4), 50 mM KCl, 1 mM DTT, 0.1 mM EDTA, 200 μ g/ml BSA, 50% (v/v) glycerol.

Heat Inactivation Condition

65°C for 20min.

Enzyme dilution

Dilute with Storage Buffer.

QC Tests

Activity, ligation and recutting, exo and endonuclease activity test, SDS-PAGE purity, performance tests.

Methylase Sensitivity

Blocked by CpG methylation (Cleavage of mammalian genomic DNA is blocked).

Buf4	37°C	65°C	CpG	BW
<small>BUFFER</small>	<small>REACTION</small>	<small>INACTIVATION</small>	<small>METHYL BLOCK</small>	<small>BLUE/WHITE</small>

Buffer	1	2	3	4
Activity	0	50	50	100