




For research use only

ISO9001

Alu I

Product	Quantity	Cat. No.	Remarks
Alu I	500 unit	EBR-1011	10 unit/ μ l

Recognition site**Source***Arthrobacter luteus.***Reaction Conditions**

Buffer 4. 20 mM Tris-acetate (pH 7.9), 50 mM potassium acetate, 10 mM magnesium acetate, 1 mM DTT. Incubate at 37°C.

Concentration & Storage Condition

Store at -20°C. 10 unit/ μ l in 10 mM Tris-HCl (pH 7.5), 100 mM KCl, 1 mM DTT, 0.1 mM EDTA, 200 μ g/ml BSA, 50% (v/v) glycerol.

Heat Inactivation Condition

65°C for 20min.

Enzyme dilution

Dilute with Storage Buffer.

QC Tests

Activity, ligation and recutting, exo and endonuclease activity test, SDS-PAGE purity, performance tests.

Methylase SensitivityNot sensitive to *dam*, *dcm* or CpG methylation.

Buf4	37°C	65°C		
<small>BUFFER</small>	<small>REACTION</small>	<small>INACTIVATION</small>	<small>METHYL BLOCK</small>	<small>BLUE/WHITE</small>

Buffer	1	2	3	4
Activity	100	100	75	100




For research use only

ISO9001

Alu I

Product	Quantity	Cat. No.	Remarks
Alu I	500 unit	EBR-1011	10 unit/ μ l

Recognition site**Source***Arthrobacter luteus.***Reaction Conditions**

Buffer 4. 20 mM Tris-acetate (pH 7.9), 50 mM potassium acetate, 10 mM magnesium acetate, 1 mM DTT. Incubate at 37°C.

Concentration & Storage Condition

Store at -20°C. 10 unit/ μ l in 10 mM Tris-HCl (pH 7.5), 100 mM KCl, 1 mM DTT, 0.1 mM EDTA, 200 μ g/ml BSA, 50% (v/v) glycerol.

Heat Inactivation Condition

65°C for 20min.

Enzyme dilution

Dilute with Storage Buffer.

QC Tests

Activity, ligation and recutting, exo and endonuclease activity test, SDS-PAGE purity, performance tests.

Methylase SensitivityNot sensitive to *dam*, *dcm* or CpG methylation.

Buf4	37°C	65°C		
<small>BUFFER</small>	<small>REACTION</small>	<small>INACTIVATION</small>	<small>METHYL BLOCK</small>	<small>BLUE/WHITE</small>

Buffer	1	2	3	4
Activity	100	100	75	100